

Upper Bounds on $K \rightarrow \pi\nu\bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$ from ε'/ε and $K_L \rightarrow \mu^+ \mu^-$ *

Andrzej J. Buras and Luca Silvestrini

*Technische Universität München, Physik Department
D-85748 Garching, Germany*

Abstract

We analyze rare kaon decays in models in which the dominant new effect is an enhanced $\bar{s}dZ$ vertex Z_{ds} . We point out that in spite of large theoretical uncertainties the CP-violating ratio ε'/ε provides at present the strongest constraint on $\text{Im } Z_{ds}$. Assuming $0 \leq \varepsilon'/\varepsilon \leq 2 \cdot 10^{-3}$ and Standard Model values for the CKM parameters we obtain the bounds $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \leq 2.4 \cdot 10^{-10}$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-) \leq 3.6 \cdot 10^{-11}$ (which are substantially stronger than the bounds found recently by Colangelo and Isidori, using ε_K instead of ε'/ε). We illustrate how these bounds can be improved with the help of the forthcoming data on ε'/ε . Using the bound on $\text{Re } Z_{ds}$ from $K_L \rightarrow \mu^+ \mu^-$ we find $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \leq 2.3 \cdot 10^{-10}$. In this context we derive an analytic upper bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ as a function of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and the short distance contribution to $\text{BR}(K_L \rightarrow \mu^+ \mu^-)$. We also discuss new physics scenarios in which in addition to an enhanced $\bar{s}dZ$ vertex also neutral meson mixing receives important new contributions. In this case larger values of the branching ratios in question cannot be excluded.

*Supported by the German Bundesministerium für Bildung und Forschung under contract 06 TM 874 and by the DFG project Li 519/2-2.

1 Introduction

Flavour-Changing Neutral Current (FCNC) processes provide a powerful tool for the tests of the Standard Model and the physics beyond it. Of particular interest are the rare kaon decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$ which are governed by Z -penguin diagrams. Within the Standard Model the branching ratios for these decays are known including next-to-leading order corrections [1, 2]. Updating the analysis in [3] we find

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = (3.1 \pm 1.3) \cdot 10^{-11}, \quad (1)$$

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (8.2 \pm 3.2) \cdot 10^{-11}, \quad (2)$$

$$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}} = (4.9 \pm 2.1) \cdot 10^{-12}, \quad (3)$$

where the errors come dominantly from the uncertainties in the CKM parameters. The corresponding theoretical uncertainties in (1)-(3) amount to at most a few percents. The reduction of $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ and the increase of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)$ with respect to the values given in [3] originate primarily from the improved experimental lower bound on the B_s^0 - $\bar{B}_s^0$ mixing and the slight increase of $|V_{ub}|$ [4]. The branching ratio in eq. (3) represents the so-called direct CP-violating contribution to $K_L \rightarrow \pi^0 e^+ e^-$. The remaining two contributions to this decay, the CP-conserving one and the indirect CP-violating one, are plagued by theoretical uncertainties [5]. They are expected to be $\mathcal{O}(10^{-12})$ but generally smaller than $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}}$. This implies that within the Standard Model $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)$ is expected to be at most 10^{-11} .

Experimentally we have [6]

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (4.2^{+9.7}_{-3.5}) \cdot 10^{-10} \quad (4)$$

and the bounds [7, 8]

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) < 1.6 \cdot 10^{-6}, \quad \text{BR}(K_L \rightarrow \pi^0 e^+ e^-) < 4.3 \cdot 10^{-9}. \quad (5)$$

Moreover from (4) and isospin symmetry one has [9] $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) < 6.1 \cdot 10^{-9}$. The data on these three branching ratios should improve considerably in the coming years.

In this context a very interesting claim has been recently made by Colangelo and Isidori [10], who analyzing rare kaon decays in supersymmetric theories pointed out

a possible large enhancement of the effective $\bar{s}dZ$ vertex leading to branching ratios as high as

$$\begin{aligned}\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) &= 4 \cdot 10^{-9}, \\ \text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) &= 1 \cdot 10^{-9}, \\ \text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}} &= 6 \cdot 10^{-10}.\end{aligned}\tag{6}$$

This amounts to an enhancement of $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ by one order of magnitude and of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)$ by two orders of magnitude relative to the Standard Model expectations. According to [10] such enhancements are still compatible with data for other FCNC processes such as $K_L \rightarrow \mu^+ \mu^-$, the K_L - K_S mass difference ΔM_K and the indirect CP violation in $K_L \rightarrow \pi\pi$ represented by the parameter ε_K . Not surprisingly these results brought a lot of excitement among experimentalists.

In this paper we would like to point out that in models in which the dominant new effect is an enhanced $\bar{s}dZ$ vertex such large enhancements of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)$ are already excluded by the existing data on the CP-violating ratio ε'/ε in spite of the large theoretical uncertainties. Similarly the large enhancement of $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ can be excluded by the data on ε'/ε and in particular by the present information on the short distance contribution to $K_L \rightarrow \mu^+ \mu^-$. The latter can be bounded by analysing the data on $\text{BR}(K_L \rightarrow \mu^+ \mu^-)$ in conjunction with improved estimates of long distance dispersive contributions [11, 12].

Our main point is as follows. Similarly to the rare decays in question also ε'/ε depends sensitively on the size of Z -penguin contributions and generally on the size of the effective $\bar{s}dZ$ vertex. In the Standard Model Z -penguins dominate the so-called electroweak penguin contributions to ε'/ε which enter this ratio with the opposite sign to QCD penguins and suppress considerably ε'/ε for large m_t [13]. If the new physics beyond the Standard Model does not modify significantly the QCD penguin contributions but enhances considerably the Z -penguin contribution, then the strong cancellation between QCD and electroweak penguin contributions to ε'/ε present in the Standard Model does not take place. Consequently large positive or negative values of ε'/ε (dependently on the sign of new contributions) can be found. While precise predictions for ε'/ε cannot be made due to substantial hadronic uncertainties, the very fact that the cancellation in question is removed makes the estimates more

reliable. Assuming that the usual analysis of the unitarity triangle involving neutral meson mixing is only weakly modified by new physics contributions we find that the values in (6) simultaneously imply $|\varepsilon'/\varepsilon| > 5 \times 10^{-3}$. This is by many standard deviations higher than the present average $(1.5 \pm 0.8) \times 10^{-3}$ extracted [14] from the experiments at CERN [15] and Fermilab [16]. As we will demonstrate below, if $0 \leq (\varepsilon'/\varepsilon)_{\text{exp}} \leq 2 \cdot 10^{-3}$ then

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \leq 2.4 \cdot 10^{-10} \quad \text{and} \quad \text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}} \leq 3.6 \cdot 10^{-11}, \quad (7)$$

which is a factor of 20 smaller than what anticipated in [10]. The new round of ε'/ε measurements could improve these bounds considerably. In this context one should note that ε'/ε is linear in the imaginary part of the $\bar{s}dZ$ vertex, whereas $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}}$ are quadratically dependent on this vertex. Thus a modest reduction of the possible range for $(\varepsilon'/\varepsilon)_{\text{exp}}$ would result in a substantial improvement of the bounds in (7).

The bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ is governed dominantly by the bound on $\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}}$ as we will explicitly demonstrate below. Reanalyzing this bound we find that the upper bound on the real part of the $\bar{s}dZ$ vertex has been overestimated in [10] by roughly a factor of two. Combining this finding with the analysis of ε'/ε we find

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \leq 2.3 \cdot 10^{-10} \quad (8)$$

which is roughly a factor of four lower than given in [10]. We would like to stress that further improvements on $(\varepsilon'/\varepsilon)_{\text{exp}}$ will have only a minor impact on (8). On the other hand improved data on $\text{BR}(K_L \rightarrow \mu^+ \mu^-)$ and in particular further improvements in the theoretical analysis of the long distance dispersive contribution to $K_L \rightarrow \mu^+ \mu^-$ could have an important impact on the bound in (8).

We also analyze a scenario of new physics in which there are substantial new contributions to neutral meson mixing so that the usual analysis of the unitarity triangle does not apply. Dependent on the sign of the new contributions the bounds on rare decays become stronger or weaker. The largest branching ratios are found when neutral meson mixing is dominated by new physics contributions that are twice as large and have opposite sign with respect to the Standard Model. This possibility is quite remote. However, if this situation could be realized in some exotic model,

then the branching ratios in question could be as high as

$$\begin{aligned}\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) &= 1.4 \cdot 10^{-9}, \\ \text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) &= 5.3 \cdot 10^{-10}, \\ \text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}} &= 2.2 \cdot 10^{-10},\end{aligned}\tag{9}$$

which are still lower than given in (6). On the other hand if the CKM matrix is assumed to be real we find the maximal values to be

$$\begin{aligned}\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) &= 3.7 \cdot 10^{-10}, \\ \text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) &= 2.9 \cdot 10^{-10}, \\ \text{BR}(K_L \rightarrow \pi^0 e^+ e^-)_{\text{dir}} &= 6.6 \cdot 10^{-11}.\end{aligned}\tag{10}$$

The paper is organized as follows. In Section 2 we summarize our strategy and present the basic formulae. In Section 3 we derive the bounds on the effective $\bar{s}dZ$ vertex using $K_L \rightarrow \mu^+ \mu^-$ and ε'/ε . In this context we derive also an analytic upper bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ as a function of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and the short distance contribution to $\text{BR}(K_L \rightarrow \mu^+ \mu^-)$. Subsequently we present implications of these bounds for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$. In Section 4 we summarize briefly our main findings.

2 Basic Formulae

We are interested in the one-loop flavour-changing effective coupling of the Z -boson to down-type quarks, in the limit of vanishing external masses and momenta. The corresponding effective Lagrangian can be generally written as

$$\mathcal{L}_{\text{FC}}^Z = \frac{G_F}{\sqrt{2}} \frac{e}{2\pi^2} M_Z^2 \frac{\cos \Theta_W}{\sin \Theta_W} Z_{ds} \bar{s} \gamma_\mu (1 - \gamma_5) d Z^\mu + \text{h.c.}\tag{11}$$

where Z_{ds} is a complex coupling. In the Standard Model one has

$$Z_{ds}^{\text{SM}} = \lambda_t C_0(x_t), \quad x_t = \frac{\overline{m}_t^2}{M_W^2},\tag{12}$$

where $\lambda_t = V_{ts}^* V_{td}$ with V_{ij} being the CKM matrix elements. $C_0(x_t)$ is a real function which for the central value of the top quark mass, $\overline{m}_t(m_t) = 166$ GeV, equals 0.79. Its explicit expression can be found in [3].

From the standard analysis of the unitarity triangle, we find

$$\text{Im } \lambda_t = (1.38 \pm 0.33) \cdot 10^{-4}, \quad \text{Re } \lambda_t = -(3.2 \pm 0.9) \cdot 10^{-4}, \quad (13)$$

and consequently

$$\text{Im } Z_{ds}^{\text{SM}} = (1.09 \pm 0.26) \cdot 10^{-4}, \quad \text{Re } Z_{ds}^{\text{SM}} = -(2.54 \pm 0.71) \cdot 10^{-4}, \quad (14)$$

where the error in m_t has been neglected.

For completeness we give the relation of Z_{ds} to U_{ds} defined by

$$\mathcal{L}_{\text{tree}}^Z = \frac{g_2}{4 \cos \Theta_W} U_{ds} \bar{s} \gamma_\mu (1 - \gamma_5) d Z^\mu + \text{h.c.} \quad (15)$$

and used in the extensions of the Standard Model in which tree level flavour-changing Z^0 couplings appear [17, 18, 19]. One has

$$U_{ds} = \frac{\sqrt{2} G_F M_W^2}{\pi^2} Z_{ds} = 1.08 \cdot 10^{-2} Z_{ds}. \quad (16)$$

Our effective coupling Z_{ds} includes both Standard Model and new physics contributions. This definition is convenient for our analysis, since it allows us to take automatically into account possible interference effects between the Standard Model and new physics. Notice however that the coupling W_{ds} used in ref. [10], and in particular the bounds derived in [10] from $K_L \rightarrow \mu^+ \mu^-$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and ε_K , do not include the Standard Model contribution. Care must be therefore taken in comparing our bounds on Z_{ds} with the bounds on W_{ds} of ref. [10].

In obtaining the formulae listed below we have used the following strategy. In the case of $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $K_L \rightarrow \pi^0 e^+ e^-$, $(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}}$ and ε'/ε we have taken the Standard Model expressions and replaced there $\lambda_t C_0$ by Z_{ds} . The remaining contributions resulting from box diagrams, gluon-penguin and photon-penguin diagrams have been evaluated for $\overline{m}_t(m_t) = 166$ GeV. That is we assume that new physics will have at most a minor impact on the latter contributions. This is for example the case in supersymmetric extensions of the Standard Model. As pointed out in ref. [10], in general supersymmetric models there is the interesting possibility of a substantial enhancement of the effective $\bar{s}dZ$ vertex with respect to its Standard Model value, via a double helicity-flipping flavour-changing squark mass insertion. On the other hand, the effect of all other supersymmetric contributions to $K \rightarrow \pi \nu \bar{\nu}$ decays via penguin and box diagrams can at most be of the order of

magnitude of the Standard Model contribution. Details can be found in ref. [20]. Furthermore, it has been shown that in general supersymmetric models the gluonic penguin contributions to ε'/ε are small due to the constraints coming from ΔM_K and ε_K , and to the negative interference between $\Delta S = 1$ box and penguin diagrams [21]. The same applies to photonic penguins. We can therefore conclude that the only place where supersymmetry can produce order-of-magnitude enhancements in the above-mentioned processes is an enhanced $\bar{s}dZ$ vertex via the Colangelo-Isidori mechanism.

We also assume that no new operators in addition to those present in the Standard Model contribute. In this case the replacement $\lambda_t C_0 \rightarrow Z_{ds}$ is justified without the modification of QCD renormalization group effects evaluated at NLO level for scales below $\mathcal{O}(m_t)$.

It should be remarked that B_0 and C_0 depend on the gauge parameter in the W-propagator. This gauge dependence cancels when both are taken into account. In using Z_{ds} instead of C_0 we assume that the latter function is hidden in Z_{ds} so that the gauge dependence in question is absent in the final formulae for branching ratios. However, it is present in Z_{ds} . The values for Z_{ds} quoted in this paper correspond then to 't Hooft-Feynman gauge. In any case, this gauge dependence is rather weak as it originates in terms which are non-leading in m_t . As box diagrams contributing to rare decays receive only small contributions from physics beyond the Standard Model we expect that the gauge dependence of new contributions to Z_{ds} is also very weak.

In deriving the formula for ε'/ε we use the NLO analytic formula for this ratio [22] which has been updated in [3]. Since ε'/ε depends visibly on $\Lambda_{\overline{MS}}^{(4)}$ we identify those coefficients in the formula in question which carry the dominant $\Lambda_{\overline{MS}}^{(4)}$ dependence. We evaluate the remaining ones for $\Lambda_{\overline{MS}}^{(4)} = 325$ MeV, corresponding to $\alpha_s^{\overline{MS}}(M_Z) = 0.118$. This leaves us first with two coefficients $r_0^{(6)}$ and $r_Z^{(8)}$ which in the range $0.113 \leq \alpha_s^{\overline{MS}}(M_Z) \leq 0.123$ are related within 10% by $r_0^{(6)} \simeq 1.1|r_Z^{(8)}|$. We will use this relation in what follows. The resulting formula for ε'/ε reproduces in the limit $Z_{ds} \rightarrow Z_{ds}^{\text{SM}}$ the Standard Model results within 10% accuracy, which is clearly sufficient for our purposes.

Our basic formulae read then as follows:

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = 1.55 \cdot 10^{-4} \left[\left(\text{Im } Z_{ds} - 4B_0 \text{Im } \lambda_t \right)^2 \right]$$

$$+\left(\text{Re } Z_{ds} + \Delta_c - 4B_0 \text{Re } \lambda_t\right)^2\Big], \quad (17)$$

where

$$\Delta_c = -(2.11 \pm 0.30) \cdot 10^{-4} \quad (18)$$

represents the internal charm contribution [1] and $B_0 = -0.182$ is the box diagram function evaluated at $\overline{m}_t(m_t) = 166$ GeV. New physics contributions are expected to arise at a scale $\geq M_W$ and are therefore included in Z_{ds} , while the charm contribution, generated at the scale m_c , can be safely evaluated in the Standard Model. Next

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = 6.78 \cdot 10^{-4} [\text{Im } Z_{ds} - 4B_0 \text{Im } \lambda_t]^2, \quad (19)$$

$$\begin{aligned} \text{BR}(K_L \rightarrow \pi^0 e^+ e^-) &= 1.19 \cdot 10^{-4} \left[\left(\text{Im } Z_{ds} - B_0 \text{Im } \lambda_t \right)^2 \right. \\ &\quad \left. + \left(\text{Im } \lambda_t + 0.08 \text{Im } Z_{ds} \right)^2 \right], \end{aligned} \quad (20)$$

$$\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}} = 6.32 \cdot 10^{-3} [\text{Re } Z_{ds} - B_0 \text{Re } \lambda_t + \bar{\Delta}_c]^2, \quad (21)$$

where

$$\bar{\Delta}_c = -(6.54 \pm 0.60) \cdot 10^{-5}. \quad (22)$$

represents the charm contribution [1].

Using (17), (19) and (21) we derive the following useful formula

$$\begin{aligned} \text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) &= 1.55 \cdot 10^{-4} \left[\pm 3.97 \sqrt{\kappa} \cdot 10^{-4} - 3B_0 \text{Re } \lambda_t + \hat{\Delta}_c \right]^2 \\ &\quad + 0.229 \cdot \text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}), \end{aligned} \quad (23)$$

where

$$\hat{\Delta} = \Delta_c - \bar{\Delta}_c = -(1.46 \pm 0.30) \cdot 10^{-4} \quad (24)$$

and κ is defined through

$$\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}} = \kappa \cdot 10^{-9}. \quad (25)$$

In evaluating $\hat{\Delta}_c$ we have included the correlation between Δ_c and $\bar{\Delta}_c$ due to their simultaneous dependence on $\Lambda_{\overline{MS}}^{(4)}$ and m_c [1]. Next we have

$$\Delta M_K = (\Delta M_K)^{\text{SM}} + 8.2 \cdot 10^{-12} \eta_{QCD} \hat{B}_K \left[(\text{Re } Z_{ds})^2 - (\text{Im } Z_{ds})^2 \right], \quad (26)$$

$$\varepsilon_K = \varepsilon_K^{\text{SM}} - 1.65 \cdot 10^3 (\text{Re } Z_{ds}) (\text{Im } Z_{ds}) \eta_{QCD} \hat{B}_K e^{i\pi/4}. \quad (27)$$

Here $(\Delta M_K)^{\text{SM}}$ and $\varepsilon_K^{\text{SM}}$ are the Standard Model box contributions and the second terms in (26) and (27) stand for Z^0 contributions. η_{QCD} is the QCD factor and $\hat{B}_K$ is a hadronic parameter. One has $\eta_{QCD}\hat{B}_K \approx 0.5$.

In the limit $\eta_{QCD}\hat{B}_K = 1$ the Z^0 contributions given here are larger by a factor of two relatively to the ones presented in [10, 17]. They agree on the other hand with those presented in [19]. For our purposes, however, this factor is irrelevant since ΔM_K and ε_K provide much weaker constraints on the coupling Z_{ds} than $K_L \rightarrow \mu^+ \mu^-$ and ε'/ε .

Finally we decompose ε'/ε as follows:

$$\frac{\varepsilon'}{\varepsilon} = \left(\frac{\varepsilon'}{\varepsilon}\right)_Z + \left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{Rest}} \quad (28)$$

and proceeding as explained above we find

$$\begin{aligned} \left(\frac{\varepsilon'}{\varepsilon}\right)_Z &= \text{Im } Z_{ds} [1.2 - R_s |r_Z^{(8)}| B_8^{(3/2)}], \\ \left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{Rest}} &= \text{Im } \lambda_t \left[-2.3 + R_s [1.1 |r_Z^{(8)}| B_6^{(1/2)} + (1.0 + 0.12 |r_Z^{(8)}|) B_8^{(3/2)}] \right]. \end{aligned} \quad (29)$$

Here

$$R_s = \left[\frac{158 \text{MeV}}{m_s(m_c) + m_d(m_c)} \right]^2. \quad (30)$$

$B_6^{(1/2)}$ and $B_8^{(3/2)}$ are non-perturbative parameters describing the hadronic matrix elements of the dominant QCD-penguin and electroweak penguin operators respectively. Finally $|r_Z^{(8)}|$ is a calculable renormalization scheme independent parameter in the analytic formula for ε'/ε in [3] which increases with $\alpha_s^{\overline{MS}}(M_Z)$ and in the range $0.113 \leq \alpha_s^{\overline{MS}}(M_Z) \leq 0.123$ takes the values

$$6.5 \leq |r_Z^{(8)}| \leq 8.5. \quad (31)$$

For R_s we will use the range

$$1 \leq R_s \leq 2, \quad (32)$$

which is compatible with the most recent lattice and QCD sum rules calculations as reviewed in [23]. We consider the ranges in (31) and (32) as conservative. Similarly we will use

$$0.8 \leq B_6^{(1/2)} \leq 1.2, \quad 0.6 \leq B_8^{(3/2)} \leq 1.0, \quad (33)$$

which is compatible with the recent lattice and large N calculations as reviewed in [3, 23]. Our treatment of $\text{Im } \lambda_t$ and $\text{Re } \lambda_t$ will be explained below.

3 Bounds from $K_L \rightarrow \mu^+ \mu^-$ and ε'/ε

3.1 Generalities

In deriving the bounds on $\text{Re } Z_{ds}$ and $\text{Im } Z_{ds}$ from $K_L \rightarrow \mu^+ \mu^-$ and ε'/ε we have to investigate whether λ_t extracted from the standard analysis of the unitarity triangle and given in (13) is significantly modified by new physics contributions. After the improved lower bound on $B_s^0 - \bar{B}_s^0$ mixing [4], the determination of λ_t in the standard analysis is governed by the values of $|V_{ub}|$, $|V_{cb}|$, ε_K and the ratio of $B_d^0 - \bar{B}_d^0$ and $B_s^0 - \bar{B}_s^0$ mixings, represented by $(\Delta M)_d/(\Delta M)_s$. The values of $|V_{ub}|$ and $|V_{cb}|$ are expected to be unaffected by new physics contributions. On the other hand, new physics can easily affect neutral meson mixings. We can therefore distinguish between three different scenarios:

- A) New physics only manifests itself through an enhanced $\bar{s}dZ$ vertex. In this case, the bounds on $\text{Re } Z_{ds}$ from $K_L \rightarrow \mu^+ \mu^-$ and on $\text{Im } Z_{ds}$ from ε'/ε imply that the Z^0 contribution to ΔM_K is fully negligible and the corresponding contribution to ε_K is below 15% of the experimental value. Consequently, the standard analysis of the unitarity triangle is only marginally affected by new physics and we can use for λ_t the values given in eq. (13).
- B) New physics affects the $\Delta F = 2$ amplitudes through some mechanism other than Z^0 -exchange (for example, this is what happens in general SUSY models such as the one considered by Colangelo and Isidori). In this case the standard analysis of the unitarity triangle is invalid. In order to be as general as possible, in this scenario we only assume unitarity of the CKM matrix. In this case

$$\text{Im } \lambda_t = |V_{ub}| |V_{cb}| \sin \delta, \quad (34)$$

with $0 \leq \delta \leq 2\pi$ and [24]

$$\left(\lambda - \frac{\lambda^3}{2} \right) |V_{cb}|_{\min}^2 (1 - R_b^{\max}) \leq -\text{Re } \lambda_t \leq \left(\lambda - \frac{\lambda^3}{2} \right) |V_{cb}|_{\max}^2 (1 + R_b^{\max}) \quad (35)$$

where $\lambda = 0.22$ is the Wolfenstein parameter and

$$R_b = \left(1 - \frac{\lambda^2}{2} \right) \frac{1}{\lambda} \left| \frac{V_{ub}}{V_{cb}} \right|. \quad (36)$$

C) Scenario B with $\text{Im } \lambda_t = 0$. Here CP-violation originates entirely in new physics contributions.

In the following two subsections we discuss the constraints on Z_{ds} from $K_L \rightarrow \mu^+ \mu^-$ and ε'/ε in scenario A, and their implications for rare K decays. Subsequently we will discuss briefly scenarios B and C.

3.2 $K_L \rightarrow \mu^+ \mu^-$ and the Bounds on $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $\text{Re } Z_{ds}$

The $K_L \rightarrow \mu^+ \mu^-$ branching ratio can be decomposed generally as follows:

$$\text{BR}(K_L \rightarrow \mu^+ \mu^-) = |\text{Re } A|^2 + |\text{Im } A|^2, \quad (37)$$

where $\text{Re } A$ denotes the dispersive contribution and $\text{Im } A$ the absorptive one. The latter contribution can be determined in a model independent way from the $K_L \rightarrow \gamma \gamma$ branching ratio. The resulting $|\text{Im } A|^2$ is very close to the experimental branching ratio $\text{BR}(K_L \rightarrow \mu^+ \mu^-) = (7.2 \pm 0.5) \cdot 10^{-9}$ [25] so that $|\text{Re } A|^2$ is substantially smaller and extracted to be [25]

$$|\text{Re } A_{\text{exp}}|^2 < 5.6 \cdot 10^{-10} \quad (90\% \text{ C.L.}). \quad (38)$$

Now $\text{Re } A$ can be decomposed as

$$\text{Re } A = \text{Re } A_{\text{LD}} + \text{Re } A_{\text{SD}}, \quad (39)$$

with

$$|\text{Re } A_{\text{SD}}|^2 \equiv \text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}} \quad (40)$$

representing the short-distance contribution given in (21). An improved estimate of the long-distance contribution $\text{Re } A_{\text{LD}}$ has been recently presented by D'Ambrosio, Isidori and Portolés [11] who find

$$|\text{Re } A_{\text{LD}}| < 2.9 \cdot 10^{-5} \quad (90\% \text{ C.L.}). \quad (41)$$

The highest possible value for $\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}}$ is found if $\text{Re } A_{\text{SD}}$ and $\text{Re } A_{\text{LD}}$ have opposite sign. The bounds (38) and (41) give then

$$\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}} < 2.8 \cdot 10^{-9}. \quad (42)$$

This result is very close to the one presented very recently by Gomez Dumm and Pich [12]. The bound in (42) should be compared with the short distance contribution within the Standard Model for which we find

$$\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}}^{\text{SM}} = (9.4 \pm 4.1) \cdot 10^{-10}. \quad (43)$$

Due to the presence of the charm contribution, the constraint on $\text{Re } Z_{ds}$ that one can extract from (42) depends on the sign of $\text{Re } Z_{ds}$. From (21) we get

$$-5.6 \cdot 10^{-4} \leq \text{Re } Z_{ds} \leq 8.1 \cdot 10^{-4}. \quad (44)$$

The upper bound is obtained using $\text{Re } \lambda_t = -4.1 \cdot 10^{-4}$ and the lower one using $\text{Re } \lambda_t = -2.3 \cdot 10^{-4}$. Note that the term $\bar{\Delta}_c$ as well as $B_0 \text{Re } \lambda_t$ play only a minor role in obtaining (44). Setting them to zero we would find $|\text{Re } Z_{ds}| \leq 6.7 \cdot 10^{-4}$.

The bound in (44) is compatible with the one of Silverman [18] who finds $|\text{Re } Z_{ds}| \leq 8.3 \cdot 10^{-4}$. To compare with ref. [10] one has to add the Standard Model contribution, since the bound $|\text{Re } W_{ds}| \leq 2.2 \cdot 10^{-3}$ in eq. (3.10) of ref. [10] only refers to the new physics contribution. We would therefore get, neglecting the charm contribution,

$$\text{Re } W_{ds} < 6.7 \cdot 10^{-4} + |\text{Re } Z_{ds}^{\text{SM}}| \leq 10^{-3}, \quad (45)$$

which is roughly a factor of two stronger than the bound in ref. [10].*

Having the upper bound on $\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{\text{SD}}$ we can use the formula (23) to derive an upper bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$. This bound is obtained by choosing the negative sign in (23), which corresponds to $\text{Re } Z_{ds} < 0$. Setting $\text{Re } \lambda_t = -4.1 \cdot 10^{-4}$, $\hat{\Delta}_c = -1.76 \cdot 10^{-4}$ we obtain

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \leq 0.229 \cdot \text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) + 2.44 \cdot 10^{-11} [\sqrt{\kappa} + 1.01]^2. \quad (46)$$

With $\kappa = 2.8$ (see (42)) this reduces to

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \leq 0.229 \cdot \text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) + 1.76 \cdot 10^{-10}. \quad (47)$$

This formula allows then to find the upper bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ once the upper bound on $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ is known. As we will now demonstrate, the latter bound can be obtained from ε'/ε .

*This difference can be traced to a factor of two missing in eq. (9) of ref. [11], as confirmed by the authors.

3.3 Constraints from ε'/ε in scenario A

In this scenario all the effects of new physics are encoded in the effective coupling Z_{ds} . As we shall verify explicitly, in this case the new physics effects in ΔM_K and ε_K can be neglected and we can use the values for λ_t given in eq. (13).

The form of the bound for $\text{Im } Z_{ds}$ from ε'/ε depends on the sign of $\text{Im } Z_{ds}$. We consider first the case $\text{Im } Z_{ds} < 0$. Then $(\varepsilon'/\varepsilon)_Z$ in (29) is positive and adds up to $(\varepsilon'/\varepsilon)_{\text{Rest}}$ which is also positive. The upper bound on $-\text{Im } Z_{ds}$ reads then

$$-\text{Im } Z_{ds} \leq \frac{\left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{max}}^{\text{exp}} - \left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{Rest}}}{\left(R_s |r_Z^{(8)}| B_8^{(3/2)} - 1.20\right)} \quad (48)$$

The most conservative bound is found by setting $\text{Im } \lambda_t = 1.05 \cdot 10^{-4}$, $R_s = 1$, $B_6^{(1/2)} = 0.8$, $B_8^{(3/2)} = 0.6$ and $|r_Z^{(8)}| = 6.5$. This gives

$$-\text{Im } Z_{ds} \leq \frac{\left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{max}}^{\text{exp}} - 4.71 \cdot 10^{-4}}{2.70}. \quad (49)$$

Setting for instance $(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}} = 2 \cdot 10^{-3}$ we find

$$-\text{Im } Z_{ds} \leq 5.66 \cdot 10^{-4}. \quad (50)$$

If $\text{Im } Z_{ds} > 0$ the bound (48) changes to

$$\text{Im } Z_{ds} \leq \frac{\left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{Rest}} - \left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{min}}^{\text{exp}}}{\left(R_s |r_Z^{(8)}| B_8^{(3/2)} - 1.20\right)} \quad (51)$$

as for large $\text{Im } Z_{ds}$ ε'/ε becomes negative and what counts is the minimal value of $(\varepsilon'/\varepsilon)^{\text{exp}}$. The most conservative bound is obtained by setting $B_6^{(1/2)} = 1.2$, $B_8^{(3/2)} = 0.6$ and $\text{Im } \lambda_t = 1.71 \cdot 10^{-4}$, and varying $r_Z^{(8)}$ and R_s in the ranges (31) and (32). It turns out that the dependence of the bound on these two parameters is rather weak. For $(\varepsilon'/\varepsilon)_{\text{min}}^{\text{exp}} = 0$ we find

$$\text{Im } Z_{ds} \leq 4.65 \cdot 10^{-4}, \quad (52)$$

which is close to the bound in (50).

Clearly the bounds on $\text{Im } Z_{ds}$ depend on the experimental values of $(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}}$ and $(\varepsilon'/\varepsilon)_{\text{min}}^{\text{exp}}$. We illustrate this in table 1. In parentheses we show the bounds one would obtain for $B_8^{(3/2)} = 1.0$. Clearly in this case the upper bounds are much stronger. The dependence of the bounds on $B_6^{(1/2)}$ is weaker than on $B_8^{(3/2)}$. Needless to say a reduction of the uncertainties in the values of $B_6^{(1/2)}$ and $B_8^{(3/2)}$ is clearly desirable.

$(\varepsilon'/\varepsilon)_{\max}^{\text{exp}}$	$2 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
$(-\text{Im } Z_{ds})_{\max} [10^{-4}]$	5.7 (2.7)	3.8 (1.8)	2.0 (0.9)
$(\varepsilon'/\varepsilon)_{\min}^{\text{exp}}$	0	$5 \cdot 10^{-4}$	$1 \cdot 10^{-3}$
$(\text{Im } Z_{ds})_{\max} [10^{-4}]$	4.7 (2.7)	3.7 (2.3)	3.2 (2.0)

Table 1: Constraints on $\text{Im } Z_{ds}$ from ε'/ε in scenario A for different values of $(\varepsilon'/\varepsilon)_{\max}^{\text{exp}}$ and $(\varepsilon'/\varepsilon)_{\min}^{\text{exp}}$ with $B_8^{(3/2)} = 0.6$ (1.0).

Using now the constraints on $\text{Im } Z_{ds}$ in table 1 we can compute the upper bounds on the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$. Similarly using (47) we can calculate the upper bound on $K^+ \rightarrow \pi^+ \nu \bar{\nu}$. These bounds are given in tables 2 and 3 for positive and negative $\text{Im } Z_{ds}$ respectively. These tables illustrate how the improvement in the allowed range for ε'/ε , to be expected in the near future, and the increase of the minimal value of $B_8^{(3/2)}$ could improve the bounds in question. The most conservative bounds in these tables are listed in (7) and (8).

$(\varepsilon'/\varepsilon)_{\min}^{\text{exp}}$	0	$5 \cdot 10^{-4}$	$1 \cdot 10^{-3}$
$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) [10^{-10}]$	2.4 (1.0)	1.7 (0.9)	1.3 (0.7)
$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-) [10^{-11}]$	3.5 (1.5)	2.4 (1.2)	1.9 (1.0)
$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) [10^{-10}]$	2.3 (2.0)	2.1 (1.9)	2.1 (1.9)

Table 2: Upper bounds for the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$, obtained in scenario A by imposing the constraints in eq. (47) and in table 1, in the case $\text{Im } Z_{ds} > 0$ with $B_8^{(3/2)} = 0.6$ (1.0).

The contributions due to Z_{ds} to ΔM_K are always negligible, while in ε_K the effects of Z_{ds} can reach 15%, and can therefore be neglected in the standard analysis of the unitarity triangle.

3.4 Constraints from ε'/ε in scenario B

We now want to discuss the most general case in which new physics contributions to neutral meson mixing can be so large as to completely invalidate the standard

$(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}}$	$2 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})[10^{-10}]$	1.6 (0.3)	0.6 (0.1)	0.1 (0.0)
$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)[10^{-11}]$	3.6 (0.9)	1.6 (0.4)	0.5 (0.2)
$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})[10^{-10}]$	2.1 (1.8)	1.9 (1.8)	1.8 (1.8)

Table 3: Upper bounds for the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$, obtained in scenario A by imposing the constraints in eq. (47) and in table 1, in the case $\text{Im } Z_{ds} < 0$ with $B_8^{(3/2)} = 0.6$ (1.0).

analysis of the unitarity triangle. In this case, we just impose unitarity of the CKM matrix and let λ_t vary in the range

$$-2 \cdot 10^{-4} \leq \text{Im } \lambda_t \leq 2 \cdot 10^{-4}. \quad (53)$$

$$1.54 \cdot 10^{-4} \leq -\text{Re } \lambda_t \leq 5.85 \cdot 10^{-4}. \quad (54)$$

These ranges are obtained from (34)-(36) using $V_{cb} = 0.040 \pm 0.003$ and $|V_{ub}/V_{cb}| = 0.091 \pm 0.016$ [4].

For positive $\text{Im } Z_{ds}$ the upper bound on $\text{Im } Z_{ds}$ is as in (51) the only modification being $(\text{Im } \lambda_t)_{\text{max}} = 2.0 \cdot 10^{-4}$. Consequently in this case the constraints on $\text{Im } Z_{ds}$ are only slightly changed relatively to scenario A, and the upper bounds on the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$ are not dramatically modified, as one can verify from tables 4 and 5. On the contrary, for negative $\text{Im } \lambda_t$ and negative $\text{Im } Z_{ds}$ there is the possibility of a large cancellation in ε'/ε between the gluon and electroweak penguin contributions leading to a possible strongly enhanced value of $\text{Im } Z_{ds}$. The upper bound on $-\text{Im } Z_{ds}$ is again given by (49) but now the most conservative bound is obtained by setting $\text{Im } \lambda_t = -2.0 \cdot 10^{-4}$, $B_8^{(3/2)} = 0.6$, $B_6^{(1/2)} = 1.2$ and varying $r_Z^{(8)}$ and R_s in the ranges (31) and (32). As shown in table 4 the maximal values of $-\text{Im } Z_{ds}$ are this time substantially larger than in scenario A and large enhancements of $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$ are possible as shown in table 6. Of course, this possibility seems quite remote, since it would correspond to the case in which neutral meson mixing is dominated by new physics contributions that are twice as large and have opposite sign with respect to the Standard Model, and are uncorrelated to the new physics contributions to Z_{ds} (any correlation between the

$(\varepsilon'/\varepsilon)_{\max}^{\text{exp}}$	$2 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
$(-\text{Im } Z_{ds})_{\max} [10^{-4}]$	12.9 (6.8)	11.0 (5.9)	9.2 (4.9)
$(\varepsilon'/\varepsilon)_{\min}^{\text{exp}}$	0	$5 \cdot 10^{-4}$	$1 \cdot 10^{-3}$
$(\text{Im } Z_{ds})_{\max} [10^{-4}]$	5.4 (3.1)	4.5 (2.7)	3.9 (2.4)

Table 4: Constraints on $\text{Im } Z_{ds}$ from ε'/ε in scenario B for different values of $(\varepsilon'/\varepsilon)_{\max}^{\text{exp}}$ and $(\varepsilon'/\varepsilon)_{\min}^{\text{exp}}$, for $-2 \cdot 10^{-4} \leq \lambda_t \leq 2 \cdot 10^{-4}$ with $B_8^{(3/2)} = 0.6$ (1.0).

different contributions can only tighten our bound). However, if this situation could be realized in some exotic model, then large enhancements of the branching ratios in question would be possible. In figure 1 we plot the upper bound on $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ for negative $\text{Im } Z_{ds}$ as a function of $\text{Im } \lambda_t$, for three different values of $(\varepsilon'/\varepsilon)_{\max}^{\text{exp}}$. It is evident that as soon as one leaves the extreme case $\text{Im } \lambda_t = -2 \cdot 10^{-4}$ the enhancement of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ quickly decreases.

With $(\text{Re } \lambda_t)_{\max} = -5.85 \cdot 10^{-4}$ the bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ in (47) modifies to

$$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) \leq 0.229 \cdot \text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) + 2.08 \cdot 10^{-10}. \quad (55)$$

This time the first term on the r.h.s is comparable to the second term and a large enhancement of $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ is possible. Simultaneously the dependence on the experimental value of ε'/ε is stronger than in the case of scenario B.

The most conservative bounds in scenario B are listed in (9).

$(\varepsilon'/\varepsilon)_{\min}^{\text{exp}}$	0	$5 \cdot 10^{-4}$	$1 \cdot 10^{-3}$
$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) [10^{-10}]$	3.2 (1.4)	2.4 (1.2)	2.0 (1.0)
$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-) [10^{-11}]$	4.7 (2.1)	3.4 (1.7)	2.8 (1.5)
$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) [10^{-10}]$	2.8 (2.4)	2.6 (2.4)	2.5 (2.3)

Table 5: Upper bounds for the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$, obtained in scenario B by imposing the constraints in eq. (55) and in table 4, in the case $\text{Im } Z_{ds} > 0$ with $B_8^{(3/2)} = 0.6$ (1.0).

$(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}}$	$2 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})[10^{-10}]$	13.9 (4.6)	10.5 (3.6)	7.6 (2.8)
$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)[10^{-11}]$	21.9 (6.9)	16.3 (5.4)	11.7 (4.0)
$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})[10^{-10}]$	5.3 (3.1)	4.5 (2.9)	3.8 (2.7)

Table 6: Upper bounds for the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$, obtained in scenario B by imposing the constraints in eq. (55) and in table 4, in the case $\text{Im } Z_{ds} < 0$ with $B_8^{(3/2)} = 0.6$ (1.0).




Figure 1: Upper bounds on $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ in scenario B as a function of $\text{Im } \lambda_t$, for negative $\text{Im } Z_{ds}$ and for three different values of $(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}}$: $(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}} = 2 \cdot 10^{-3}$ (curve I), $(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}} = 1.5 \cdot 10^{-3}$ (curve II) and $(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}} = 1 \cdot 10^{-3}$ (curve III).

3.5 Constraints from ε'/ε in scenario C

Finally we consider scenario C, which corresponds to scenario B in which the CKM matrix is real and $\text{Im } \lambda_t$ vanishes. CP-violation originates then entirely in new physics contributions. In this case we find

$$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = 6.78 \cdot 10^{-4} [\text{Im } Z_{ds}]^2 \quad (56)$$

$$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-) = 1.20 \cdot 10^{-4} [\text{Im } Z_{ds}]^2 \quad (57)$$

and the bound (55). If $(\varepsilon'/\varepsilon)^{\text{exp}}$ is positive then only $\text{Im } Z_{ds} < 0$ is allowed. Then we have

$$-\text{Im } Z_{ds} \leq \frac{\left(\frac{\varepsilon'}{\varepsilon}\right)_{\text{max}}^{\text{exp}}}{\left(6.5 B_8^{(3/2)} - 1.20\right)} \quad (58)$$

and the numerical results in table 7. This time the upper bounds on $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)$ are substantially stronger than in the case of scenario B. The most conservative bounds in scenario C are listed in (10).

$(\varepsilon'/\varepsilon)_{\text{max}}^{\text{exp}}$	$2 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$	$1 \cdot 10^{-3}$
$(-\text{Im } Z_{ds})_{\text{max}} [10^{-4}]$	7.4 (3.8)	5.6 (2.8)	3.7 (1.9)
$\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu}) [10^{-10}]$	3.7 (1.0)	2.1 (0.5)	0.9 (0.2)
$\text{BR}(K_L \rightarrow \pi^0 e^+ e^-) [10^{-11}]$	6.6 (1.7)	3.7 (1.0)	1.6 (0.4)
$\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) [10^{-10}]$	2.9 (2.3)	2.6 (2.2)	2.3 (2.1)

Table 7: Upper bounds for the rare decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 e^+ e^-$, obtained in scenario C in the case $\text{Im } Z_{ds} < 0$ with $B_8^{(3/2)} = 0.6$ (1.0).

4 Summary

In this paper we have considered the possibility of an enhanced $\bar{s}dZ$ vertex Z_{ds} . We have pointed out that in spite of large theoretical uncertainties the best present constraints on $\text{Im } Z_{ds}$ follow from ε'/ε . These constraints should be considerably improved when new data on ε'/ε will be available and the theoretical uncertainties reduced. While considerable enhancements of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow$

$\pi^0 e^+ e^-$) over the Standard Model expectations are still possible, the huge enhancements by two orders of magnitude claimed in [10] are in our opinion already excluded. This is in particular the case in scenarios in which the analysis of the unitarity triangle is only insignificantly modified by new physics contributions.

Similar comments apply to $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ which is dominantly bounded by $\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{SD}$. Using the most recent estimates of the long distance dispersive contribution to $\text{BR}(K_L \rightarrow \mu^+ \mu^-)$ [11, 12] we find that values of $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ of the order of 10^{-9} are certainly excluded. In this context we have presented an analytic upper bound on $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ as a function of $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \mu^+ \mu^-)_{SD}$.

Our results for various bounds are collected in tables 1-7 and figure 1 with the most conservative bounds listed in (7), (8), (9) and (10). Clearly the best constraints on $\text{Im } Z_{ds}$ and $\text{Re } Z_{ds}$ will follow in the future from precise measurements of the theoretically clean branching ratios $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ respectively. Meanwhile it will be exciting to follow the development in the improved experimental values for ε'/ε , $\text{BR}(K_L \rightarrow \mu^+ \mu^-)$, $\text{BR}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, $\text{BR}(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $\text{BR}(K_L \rightarrow \pi^0 e^+ e^-)$.

Acknowledgements

We warmly thank Gilberto Colangelo, Gino Isidori and Yossi Nir for discussions and clarifications about their work. We would also like to thank Gustavo Burdman, Louis Fayard, Laurie Littenberg and Bruce Winstein for asking questions related to [10], which motivated this work.

References

- [1] G. Buchalla and A.J. Buras, *Nucl. Phys. B* **400** (1993) 225; *Nucl. Phys. B* **412** (1994) 106.
- [2] A.J. Buras, M.E. Lautenbacher, M. Misiak and M. Münz, *Nucl. Phys. B* **423** (1994) 349.

- [3] A.J. Buras, hep-ph/9806471, to appear in *Probing the Standard Model of Particle Interactions*, eds. F. David and R. Gupta (Elsevier Science B.V., Amsterdam, 1998).
- [4] J. Alexander; D. Karlen; P. Rosnet; talks given at the 29th International Conference on High Energy Physics (ICHEP 98), Vancouver, Canada, 23-29 July 1998, to appear in the proceedings.
- [5] G. Ecker, A. Pich and E. de Rafael, *Nucl. Phys.* **B291** (1987) 692, *Nucl. Phys.* **B303** (1988) 665;
A.G. Cohen, G. Ecker and A. Pich, *Phys. Lett.* **B304** (1993) 347;
P. Heiliger and L. Seghal, *Phys. Rev.* **D47** (1993) 4920;
C. Bruno and J. Prades, *Z. Phys.* **C57** (1993) 585;
J.F. Donoghue and F. Gabbiani, *Phys. Rev.* **D51** (1995) 2187;
A. Pich, hep-ph/9610243.
- [6] S. Adler et al., *Phys. Rev. Lett.* **79** (1997) 2204.
- [7] J. Adams et al., hep-ex/9806007.
- [8] D.A. Harris et al., *Phys. Rev. Lett.* **71** (1993) 3918.
- [9] Y. Grossman and Y. Nir, *Phys. Lett.* **B 398** (1997) 163.
- [10] G. Colangelo and G. Isidori, *JHEP* 09 (1998) 009.
- [11] G. D'Ambrosio, G. Isidori and J. Portolés, *Phys. Lett.* **B 423** (1998) 385.
- [12] D. Gomez Dumm and A. Pich, *Phys. Rev. Lett.* **80** (1998) 4633.
- [13] J.M. Flynn and L. Randall, *Phys. Lett.* **B 224** (1989) 221;
G. Buchalla, A.J. Buras and M.K. Harlander, *Nucl. Phys.* **B 337** (1990) 313.
- [14] C. Caso et al., Review of Particle Physics, *Eur. Phys. J.* **C 3** (1998) 1.
- [15] G.D. Barr et al., *Phys. Lett.* **B 317** (1993) 233.
- [16] L.K. Gibbons et al., *Phys. Rev. Lett.* **70** (1993) 1203.
- [17] Y. Nir and D. Silverman, *Phys. Rev.* **D 42** (1990) 1477.

- [18] D. Silverman, *Phys. Rev.* **D 45** (1992) 1800.
- [19] G. Barenboim and F.J. Botella, *Phys. Lett.* **B 433** (1998) 385.
- [20] Y. Nir and M.P. Worah, *Phys. Lett.* **B 423** (1998) 319;
A.J. Buras, A. Romanino and L. Silvestrini, *Nucl. Phys.* **B 520** (1998) 3.
- [21] E. Gabrielli, A. Masiero and L. Silvestrini, *Phys. Lett.* **B 374** (1996) 80;
F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, *Nucl. Phys.* **B 477** (1996) 321;
A. Masiero and L. Silvestrini, hep-ph/9709242.
- [22] A.J. Buras and M.E. Lautenbacher, *Phys. Lett.* **B 318** (1993) 212;
A.J. Buras, M. Jamin and M.E. Lautenbacher, *Phys. Lett.* **B 389** (1996) 749.
- [23] R. Gupta, hep-ph/9801412.
- [24] A.J. Buras, M.E. Lautenbacher and G. Ostermaier, *Phys. Rev.* **D 50** (1994) 3433.
- [25] A.P. Heinson et al., *Phys. Rev.* **D 51** (1995) 985;
T. Akagi et al., *Phys. Rev.* **D 51** (1995) 2061.